

# 长期施用化肥和有机肥下潮土干团聚体有机氮组分特征\*

李 娇<sup>1,2</sup> 信秀丽<sup>1</sup> 朱安宁<sup>1†</sup> 杨文亮<sup>1</sup> 侯 滢<sup>3</sup>

邹文萱<sup>3</sup> 徐琳雅<sup>3</sup>

(1 土壤与农业可持续发展国家重点实验室/封丘农业生态实验站(中国科学院南京土壤研究所), 南京 210008)

(2 中国科学院大学, 北京 100049)

(3 南京农业大学资源与环境科学学院, 南京 210095)

**摘要** 依托中国科学院封丘农业生态实验站长期施肥试验, 选取不施肥(CK)、单施化肥(NPK)、有机无机肥配施(1/2OM)、有机肥(OM)处理, 对比研究 27 a 连续施用化肥和有机肥对土壤机械稳定性团聚体及其有机氮组分的影响。结果表明, 长期施用有机肥显著提高了耕层土壤中大于 2 mm 团聚体的比例, 较 CK 提高了 33%, 较 NPK 增加了 17%。施肥显著提高耕层团聚体中有机氮含量, 以 OM 处理效果最明显, 大于 2 mm、0.25~2 mm、小于 0.25 mm 团聚体中酸解有机氮含量分别为 776.4、837.7、625.3 mg kg<sup>-1</sup>。各团聚体中有机氮以酸解铵态氮为主, 氨基糖态氮最少。长期单施化肥主要提高了大于 2 mm 团聚体中酸解铵态氮比例, 施用有机肥提高了氨基酸态氮和酸解未知态氮含量及分配比例。长期施用有机肥使潮土结构明显改善, 有利于耕层团聚体中全氮及有机氮各组分的积累, 氨基酸态氮、氨基糖态氮、非酸解有机氮主要赋存于 0.25~2 mm 团聚体中, 而酸解铵态氮和酸解未知态氮在大于 2 mm 团聚体中分布较多, 有效地提高了土壤供氮能力。

**关键词** 长期施肥; 潮土; 团聚体; 酸解有机氮; 全氮

**中图分类号** S153.6+2 **文献标识码** A

团聚体是土壤结构的重要组成, 其形成及稳定与施肥管理密切相关<sup>[1]</sup>。刘哲等<sup>[2]</sup>研究发现有机肥施入能提高温室和农田土壤中大团聚体含量及稳定性; 施用生物肥料有利于滨海盐碱地有机碳的积累和大团聚体的形成<sup>[3]</sup>, 但也有研究表明禽畜粪便对团聚体稳定性影响较小<sup>[4]</sup>。土壤中不同粒级团聚体对氮素的吸附和保护能力存在一定差异, 进而对各形态氮素的分布和组成产生影响<sup>[5-6]</sup>。

土壤有机氮是土壤氮素的主要存在形态, 也是矿质态氮的源和库, 在土壤养分保蓄、氮素循环及粮食生产中发挥重要作用<sup>[7]</sup>, 长久以来受到国内外学者的广泛关注。Karamjit 等<sup>[8]</sup>对稻麦轮作下氮素形态研究表明, 单施化肥显著提高了土壤中全氮和有机氮含量; 李萌等<sup>[9]</sup>报道, 猪粪代替化肥能显著增加酸解有机氮含量, 由高到低依次为非酸解有机氮、氨基酸态氮、酸解铵态氮、酸解未知氮、氨基糖态氮。不同轮作模式下, 周年氮肥施用量超过 300 kg hm<sup>-2</sup>

\* 国家重点研发计划项目(2016YFD0200304、2016YFD0300809)和江苏省重点研发计划项目(BE2017343-1)资助 Supported by the National Key Research and Development Program of China (Nos.2016YFD0200304 and 2016YFD0300809) and the Key Research and Development Program of Jiangsu Province in China (No. BE2017343-1)

† 通讯作者 Corresponding author, E-mail: [anzhu@issas.ac.cn](mailto:anzhu@issas.ac.cn)

作者简介: 李 娇(1993—), 女, 河南新乡人, 硕士研究生, 主要从事土壤氮素循环相关研究。E-mail: [lijiao@issas.ac.cn](mailto:lijiao@issas.ac.cn)

收稿日期: 2018-03-01; 收到修改稿日期: 2018-06-09; 优先数字出版日期(www.cnki.net):

时,土壤全氮及有机氮各组分含量均有所增加<sup>[10]</sup>。Kaur 和 Singh<sup>[11]</sup>对黏壤土中有机氮分析发现,有机无机肥配施下酸解有机氮较单施化肥进一步提高了 7%~34%, Xu 等<sup>[12]</sup>分析了壤黏土不同颗粒中有机氮含量,发现有机无机肥配施主要增加氨基糖态氮及大于 2 μm 粒径中氨基酸态氮含量,随着有机肥施用量的增加,以小于 2 μm 粒级中酸解有机氮含量最高<sup>[13]</sup>。

黄淮海平原被誉为中国的粮仓,该地区土壤砂粒含量高、结构较差,较大程度地影响了土壤氮素的保蓄与供应能力。因此,开展长期不同施肥措施对潮土团聚体形成及其有机氮组分变化的研究,对于深入探究施肥对潮土供氮能力及氮素转化过程的影响具有重要意义。本研究基于黄淮海平原长期定位施肥试验,研究长期不同施肥措施下潮土团聚结构变化特征,明确施肥对团聚体氮素形态含量与组成的影响,为制定合理的施肥措施、提升土壤肥力提供理论支持。

## 1 材料与方法

### 1.1 试验区概况

试验在中国科学院封丘农业生态实验站(35°00'N,114°24'E)内进行。该地区主要气候类型为半干旱半湿润的暖温带季风气候,年均气温为 13.9 °C,多年平均降水量约 605 mm,无霜期 220 d 左右。土壤类型为黄河冲积物发育形成的典型潮土,耕层土壤质地为砂壤土。

### 1.2 试验设计

长期定位施肥试验始于 1989 年秋季,冬小麦/夏玉米轮作一年两熟制,试验开始前耕层土壤(0~20 cm)的理化性质为:有机质 5.83 g kg<sup>-1</sup>、全氮 0.445 g kg<sup>-1</sup>、有效磷 1.93 mg kg<sup>-1</sup>、速效钾 78.8 mg kg<sup>-1</sup>、pH 8.65,其他理化性质及试验设计参见文献[14]。为探究长期均衡施肥对土壤团聚体形成及其有机氮组分的影响,本研究选取:不施肥(CK)、化肥氮磷钾(NPK)、1/2 有机肥氮+1/2 化肥氮(1/2OM)、有机肥(OM)四个处理,肥料品种分别为:尿素、过磷酸钙、硫酸钾,施肥量见表 1。

表 1 各处理年施肥量

Table 1 Fertilizer application rate in the experiment relative to treatment/(kg hm<sup>-2</sup>)

处理 Treatment	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
CK	0	0	0
NPK	300	135	300
1/2OM	300	135	300
OM	300	135	300

注: CK: 不施肥, NPK: 化肥氮磷钾, 1/2OM: 1/2 有机肥氮+1/2 化肥氮, OM: 有机肥。下同 Note: CK: no fertilization, NPK: Application of chemical fertilizers only, 1/2OM: Application of half the rate of chemical fertilizer in Treatment NPK and half of the rate of organic manure in Treatment OM, and OM: Application of organic manure only

### 1.3 样品采集与分析

2016 年 10 月夏玉米收获后,采集 0~20 cm 土壤样品,每个小区以“S”形多点取样,将大块土样沿自然裂缝掰开,于室内避光处自然风干。采用沙维诺夫干筛法<sup>[15]</sup>对土样进行筛分,分离获得 3 种粒径的团聚体,分别为>2 mm、0.25~2 mm、<0.25 mm。

土壤有机氮分级测定采用 Bremner 法<sup>[16]</sup>,其中,酸解有机氮(Total acidolizable nitrogen)用凯氏蒸馏法测定,酸解铵态氮(Acidolizable ammonia nitrogen)用氧化镁蒸馏法测定,铵

态氮和氨基糖态氮 (Amino sugar nitrogen) 的总量用磷酸盐-硼酸盐缓冲液蒸汽蒸馏法测定, 氨基酸态氮 (Amino acid nitrogen) 用茚三酮氧化/磷酸盐-硼酸盐氧化法测定, 差减法获得非酸解性氮 (Non-acidolizable nitrogen)、氨基糖态氮、酸解未知态氮 (Unknown-acidolizable nitrogen) 含量 (见式 (1)、式 (2)、式 (3))。土壤全氮采用半微量凯氏法测定<sup>[17]</sup>。

$$\text{非酸解有机氮} = \text{全氮} - \text{酸解有机氮} \quad (1)$$

$$\text{氨基糖态氮} = [(\text{酸解铵态氮} + \text{氨基糖态氮}) - \text{酸解铵态氮}] \times 1.4 \quad (2)$$

$$\text{酸解未知态氮} = \text{酸解有机氮} - \text{酸解铵态氮} - \text{氨基酸态氮} - \text{氨基糖态氮} \quad (3)$$

#### 1.4 数据处理

供试土壤各级团聚体的质量比例及其对养分的贡献率的计算公式如下:

$$\text{各级团聚体的质量比例} = \frac{\text{各处理中该级团聚体的质量}}{\text{各处理土壤样品总质量}} \times 100\% \quad (4)$$


$$\text{团聚体贡献率} = \frac{\text{该级团聚体中养分含量} \times \text{该级团聚体的质量比例}}{\text{耕层土壤中养分含量}} \times 100\% \quad (5)$$

试验数据采用 Excel 2010 和 SPSS 21.0 进行计算和单因素方差分析, 利用最小显著差异 (LSD) 法进行显著性检验。

## 2 结果

### 2.1 不同施肥处理土壤团聚体及全氮

各处理中大于 2 mm 团聚体质量比例范围在 39%~54% (图 1), 1/2OM、OM 显著提高了大于 2 mm 团聚体质量比例 ( $P < 0.05$ ), 较 CK 分别提高了 13%~14%, 较 NPK 分别增加了 17%~18%, 而 0.25~2 mm 团聚体质量比例无显著变化。同时, 施肥使耕层土壤中小于 0.25 mm 团聚体质量比例降低 19%~37%。



注: 不同小写字母表示同一粒径在不同处理间差异达到显著水平 ( $P < 0.05$ )。下同 Note: Different lowercase letters mean significant differences ( $P < 0.05$ ) between treatments within the same fraction of aggregates. The same below

图 1 不同施肥处理下土壤团聚体的质量比例

Fig. 1 Mass proportion of aggregates relative to treatment

连续施肥 27 a 后, 耕层土壤团聚体中全氮含量显著增加 (图 2 a)。与 CK 相比, 长期单施化肥显著提高团聚体中全氮含量, 随着团聚体粒径减小, 增加率分别为 60%、68%、60%; 施用有机肥进一步促进了团聚体中全氮的积累, 且 OM 处理效果优于 1/2OM, 随着团聚体粒径减小, 较 1/2OM 处理分别增加 16%、26%、20%。各施肥处理下, 不同粒径团聚体中全氮含量由高到低依次为 0.25~2 mm、大于 2 mm、小于 0.25 mm 的趋势。

不同施肥处理中团聚体对氮素的贡献率有所不同,如图 2 b 所示。施用有机肥显著提高了大于 2 mm 团聚体对全氮的贡献率,较 CK 增加了 36%~40%,较 NPK 增加了 21%~24%。与不施肥相比,施肥降低了小于 0.25 mm 团聚体对全氮的贡献,1/2OM 处理效果最为显著。




图 2 土壤团聚体中全氮含量 (a) 及其对土壤全氮的贡献率 (b)

Fig.2 Content of total nitrogen in soil aggregates (a) and contribution to TN (b) of the soil relative to fraction of aggregates and treatment

## 2.2 不同团聚体有机氮含量及组成

各施肥处理下团聚体内酸解有机氮含量为 147.9~837.7 mg kg<sup>-1</sup>, 占土壤全氮的 36%~72% (表 2)。施肥显著增加团聚体中酸解有机氮含量, 以 OM 处理效果最佳, 较 CK 提高 4 倍 ( $P<0.05$ ), 较 NPK 提高 94% ( $P<0.05$ ), 且有机氮主要赋存于大于 0.25 mm 团聚体中。非酸解有机氮含量为 253.2~497.6 mg kg<sup>-1</sup>, 占土壤全氮的 28%~64%, 以 0.25~2 mm 团聚体中分布最多。

经过 27 a 的培肥过程, NPK 处理氨基酸态氮含量显著高于 CK, 1/2OM、OM 处理较 NPK 分别提高了 27%~52%、45%~95% (表 2)。CK 处理中小于 0.25 mm 团聚体中氨基酸态氮含量最高, 0.25~2 mm 团聚体中含量最低, NPK、1/2OM、OM 处理与之相反。团聚体中氨基酸态氮占土壤全氮的 11%~23%, 与单施化肥相比, 有机肥主要提高了大于 0.25 mm 团聚体氨基酸态氮含量及分配比例。

酸解铵态氮含量为 70.3~265.8 mg kg<sup>-1</sup>, 占全氮的 15%~24%。NPK、1/2OM、OM 与 CK 相比分别提高了 0.8 倍~1.2 倍、1.4 倍~1.9 倍、2.0 倍~2.8 倍, 1/2OM、OM 处理较 NPK 分别增加 28%~33%、57%~73%。与其他施肥处理相比, 长期单施化肥主要提高了大于 2 mm 团聚体中酸解铵态氮比例。

团聚体中氨基糖态氮含量最低(3.9~15.0 mg kg<sup>-1</sup>), 在土壤氮素中分配比例仅为 1%~2%。各施肥处理均显著提高了团聚体中氨基糖态氮含量, 以 1/2OM 处理含量最高, 占土壤全氮的 1.1%~1.4%。团聚体中氨基糖态氮含量由高到低依次为 0.25~2 mm、大于 2 mm、小于 0.25 mm。

酸解未知态氮含量范围为 14.4~271.6 mg kg<sup>-1</sup>, 占全氮的 4%~27%。与 CK 相比, NPK 处理酸解未知态氮含量显著提高 1.6 倍~5.5 倍, 以大于 0.25 mm 团聚体酸解未知态氮含量居高; 1/2OM、OM 处理较 NPK 进一步提高了 0.53 倍~1.84 倍。

表 2 不同施肥处理下团聚体有机氮各组分含量及组成

Table 2 Content and proportion of organic nitrogen in soil aggregates relative to fraction and treatment

有机氮 organic nitrogen	处理 Treatment	>2mm		0.25~2 mm		<0.25 mm	
		含量 content/(mg k g <sup>-1</sup> )	占全氮比例 Ratio to total nitrogen/%	含量 content/( mg kg <sup>-1</sup> )	占全氮比例 Ratio to total nitrogen/%	含量 content/(mg k g <sup>-1</sup> )	占全氮比例 Ratio to total nitrogen/%
酸解有机氮	CK	176.9±4.1d	41	166.3±4.1 d	36	147.9±2.2 d	37
Total	NPK	437.7±8.2 c	64	430.8±6.0 c	55	348.6±24.3 c	54
acidolyzable	1/2OM	701.7±8.0 b	72	621.2±8.0 b	59	473.0±20.5 b	58
nitrogen	OM	776.4±9.2 a	68	837.7±12.3 a	63	657.3±13.4 a	67
非酸解有机	CK	257.2±4.1 bc	59	301.5±4.1 d	64	253.2±2.2 c	63
氮	NPK	249.4±8.2 c	36	354.6±6.0 c	45	291.3±24.3 b	46
Non-acidolyza	1/2OM	270.1±13.5 b	28	435.0±8.0 b	41	347.8±20.5 a	42
ble nitrogen	OM	357.1±9.2 a	32	497.6±12.2 a	37	325.4±15.5 a	33
氨基酸态氮	CK	46.0±1.0 d	11	53.8±3.4 d	11	58.0±1.7 d	14
Amino acid	NPK	136.6±4.6 c	20	146.6±8.3 c	19	116.7±9.68 c	18
nitrogen	1/2OM	201.6±1.8 b	21	223.1±5.2 b	21	147.9±1.4 b	18
	OM	255.0±3.7 a	23	285.3±12.5 a	21	169.2±12.6 a	17
酸解铵态氮	CK	74.7±3.0 d	17	70.3±2.8 d	15	71.5±1.3 d	18
Acidolyzable	NPK	161.8±4.9 c	24	153.2±2.9 c	20	131.6±1.2 c	21
ammonia	1/2OM	214.6±4.5 b	22	202.3±0.4 b	19	168.8±5.0 b	21
nitrogen	OM	253.7±5.6 a	22	265.8±6.2 a	20	213.7±1.9 a	22
氨基糖态氮	CK	6.4±0.6 c	1	3.9±0.9 b	1	4.1±0.3 c	1
Amino sugar	NPK	11.3±1.4 b	2	12.9±0.9 a	2	6.2±1.6 b	1
nitrogen	1/2OM	14.0±0.7 a	1	14.8±2.6 a	1	9.1±0.67 a	1
	OM	11.1±0.8 b	1	15.0±2.3 a	1	6.6±0.8 b	1
酸解未知态	CK	49.9±1.7 d	11	38.3±3.0 d	8	14.4±2.5 d	4
氮	NPK	128.0±6.5 c	19	118.2±17.1 c	15	94.1±8.8 c	15
Unknown-acid	1/2OM	271.5±5.4 a	28	181.0±4.4 b	17	147.2±35.4 b	18
olyzable	OM	256.6±4.2 b	23	271.6±7.6 a	20	267.7±24.0 a	27
nitrogen							

注: 表中数据为平均值±标准差, 不同小写字母表示同一粒径团聚体质量比例在不同处理间差异达显著水平 ( $P<0.05$ ) Note: Data in the table are of mean±standard deviation. Different lowercase letters mean significant differences ( $P<0.05$ ) between different treatments within the same fraction of soil aggregates

### 3 讨论

#### 3.1 长期单施化肥对潮土团聚体及其有机氮组分的影响

土壤质量的优劣与土壤团聚体数量及组成密切相关。与不施肥相比, 长期施用化肥提高了耕层土壤中大于 2 mm 团聚体含量, 与 Chen 等<sup>[18]</sup>结果相似。作物根系及真菌生物量是影响土壤大团聚体形成的主要生物因素<sup>[19]</sup>, 化肥氮的施入有效提高作物根系和真菌数量<sup>[20]</sup>, 进而影响土壤大团聚体的形成。同时, 土壤中大于 2 mm 团聚体对全氮贡献率最高, 表明大

团聚在养分保蓄方面发挥重要作用。

与其他处理相比,长期单施化肥主要提高了大于 2 mm 团聚体中酸解铵态氮比例。酸解铵态氮有部分源自土壤中无机态的交换性铵和部分固定态铵<sup>[21]</sup>,施用化肥增加了土壤中固定态铵的含量,有利于酸解铵态氮的累积。此外,有研究表明,化肥氮对土壤大团聚体的影响更大<sup>[20]</sup>。丛耀辉<sup>[22]</sup>研究报道:在大于 2 mm 团聚体中,酸解铵态氮与该粒级团聚体组成关系密切,可能是由于大于 2 mm 团聚体中具有较多的吸附性铵和固定态铵。1/2OM、OM 处理中酸解铵态氮含量虽然增加,但是其比例低于 NPK 处理。

### 3.2 长期施用有机肥对团聚体及其有机氮组分的影响

有机质作为土壤团聚体的重要胶结物质,在其形成过程中发挥了重要作用。长期施用有机肥使土壤有机质含量大幅度提高<sup>[23]</sup>,同时增加了作物残茬的残留,有利于土壤团聚体的形成。本研究中,施用有机肥显著增加了耕层土壤大于 2 mm 团聚体比例(图 1),改善了土壤物理性质。但是,刘中良等<sup>[24]</sup>研究表明,连续 8 a 大量施用有机肥减少了 >2 mm 团聚体的分布,可能是由于以下两个方面原因:一是土壤本身性质产生的差异,本研究中所用的砂性潮土,砂粒含量高、有机质含量较低,而刘中良等<sup>[24]</sup>研究所用的潮棕壤有机质含量高达 191.8 g kg<sup>-1</sup>;另一方面是施肥量不同,土壤固定有机碳的能力有限,过量输入有机肥导致大于 2 mm 团聚体中有机碳更新速度加快,从而不利于大于 2 mm 团聚体的形成。

有机肥处理显著提高潮土耕层土壤中氨基酸态氮和酸解未知态氮含量及分配比例。氨基酸态氮主要为蛋白质分解产物,受到土壤腐殖质和无机物(如黏土矿物和铁铝氧化物)的保护<sup>[25]</sup>。施用有机肥显著增加团聚体中氨基酸态氮占全氮比例,主要是通过影响微生物“矿化-固持”改变有机氮组分结构<sup>[26]</sup>。土壤中的可矿化氮主要来自酸解有机氮,尤其是氨基酸态氮和酸解铵态氮<sup>[7,27]</sup>,施用有机肥增加了土壤中易矿化氮含量,增强土壤氮素的供应能力<sup>[28]</sup>。

### 3.3 长期有机无机肥配施对团聚体及其有机氮组分的影响

有机无机肥配施对团聚体的影响与 OM 处理相似,均促进了土壤大团聚体的形成。骆坤等<sup>[29]</sup>研究显示,有机无机配施尤其有利于提高耕层土壤有机碳氮的活性;也有研究认为有机无机肥配施有可能加速有机物质的分解,从而影响团聚作用,这可能与有机无机肥的配比有关<sup>[30]</sup>。

本研究中,有机肥无机肥配施能显著提高大团聚体中氨基糖态氮含量,且在大团聚体中的含量高于微团聚体,与 Ding 等<sup>[31]</sup>研究结果相似。主要是由于氨基糖态氮主要源于微生物细胞壁的残留物<sup>[10]</sup>,而高有机物质投入后直接增强了多数微生物活性以及提高代谢物(包括微生物细胞壁残留物)的产量<sup>[32]</sup>。徐阳春等<sup>[33]</sup>认为,肥料配施条件下,土壤氨基糖态氮在氮素循环转化过程中具有较强的稳定性;也有研究<sup>[22]</sup>表明,氨基糖态氮在由黏粒向大团聚体团聚过程中起着重要的胶结作用。总体而言,氨基糖态氮含量虽然少,但是其在土壤团聚体结构形成的过程中发挥着重要作用。

## 4 结论

长期施用有机肥显著提高了潮土耕层大于 2 mm 团聚体的比例,同时降低小于 0.25 mm 团聚体质量比例,改善了土壤结构。施肥显著影响团聚体中氮素形态及含量分布,各级团聚体中酸解有机氮组分由高到低依次为酸解铵态氮、氨基酸态氮、酸解未知态氮、氨基糖态氮。其中,长期单施化肥主要提高了大于 2 mm 团聚体中酸解铵态氮比例,施用有机肥显著增加了氨基酸态氮和酸解未知态氮含量及分配比例,可有效提高土壤氮素的供应能力。

参考文献

- [1] 陈晓芬, 李忠佩, 刘明, 等. 不同施肥处理对红壤水稻土团聚体有机碳、氮分布和微生物生物量的影响. 中国农业科学, 2013, 46(5): 950-960  
Chen X F, Li Z P, Liu M, et al. Effects of different fertilizations on organic carbon and nitrogen contents in water-stable aggregates and microbial biomass content in paddy soil of subtropical China (In Chinese). Scientia Agricultura Sinica, 2013, 46(5): 950-960
- [2] 刘哲, 孙增慧, 吕贻忠. 长期不同施肥方式对华北地区温室和农田土壤团聚体形成特征的影响. 中国生态农业学报, 2017, 25(8): 1119-1128  
Liu Z, Sun Z H, Lü Y Z. Effects of long-term fertilization on soil aggregates formation in greenhouse and in farmland in Huabei Plain, China (In Chinese). Chinese Journal of Eco-Agriculture, 2017, 25(8): 1119-1128
- [3] Cong P P, Zhu O Y, Hou R X et al. Effects of application of microbial fertilizer on aggregation and aggregate-associated carbon in saline soils. Soil and Tillage Research, 2017, 168: 33-41
- [4] Bottinelli N, Angers D A, Hallaire V. et al. Tillage and fertilization practices affect soil aggregate stability in a Humic Cambisol of Northwest France. Soil and Tillage Research, 2017, 170: 14-17
- [5] Kumari M, Chakraborty D, Gathala M K. et al. Soil aggregation and associated organic carbon fractions as affected by tillage in a rice-wheat rotation in North India. Soil Science Society of America Journal, 2011, 75(2): 560-567
- [6] Hao Y J, Wang Y H, Chang Q R, et al. Effects of long-term fertilization on soil organic carbon and nitrogen in a highland agroecosystem. Pedosphere, 2017, 27(4): 725-736
- [7] Li S X, Wang Z H, Miao Y F, et al. Soil organic nitrogen and its contribution to crop production. Journal of Integrative Agriculture, 2014, 13(10): 2061-2080
- [8] Karamjit S S, Jai P S, Dalel S M. Long-term effect of manure and mineral fertilizer application on the distribution of organic nitrogen fractions in soil under a rice-wheat cropping system. Archives of Agronomy and Soil Science, 2011, 57(7): 705-714
- [9] 李萌, 王昌全, 李冰, 等. 猪粪替代氮肥对稻麦轮作条件下土壤有机氮组分的影响. 土壤, 2016, 48(3): 449-454  
Li M, Wang C Q, Li B, et al. Effects of pig manure replacing nitrogen fertilizer on soil organic nitrogen components under rice-wheat rotation (In Chinese). Soils, 2016, 48(3): 449-454
- [10] 贾倩, 廖世鹏, 卜容燕, 等. 不同轮作模式下氮肥用量对土壤有机氮组分的影响. 土壤学报, 2017, 54(6): 1547-1558  
Jia Q, Liao S P, Bo R Y, et al. Effects of nitrogen application rate on fractionation of soil organic nitrogen relative to crop rotation mode (In Chinese). Acta Pedologica Sinica, 2017, 54(6): 1547-1558
- [11] Kaur J, Singh J P. Long-term effects of continuous cropping and different nutrient management practices on the distribution of organic nitrogen in soil under rice-wheat system. Plant, Soil and Environment, 2014, 60(2): 63-68
- [12] Xu Y C, Shen Q R, Ran W. Content and distribution of forms of organic N in soil and particle size fractions after long-term fertilization. Chemosphere, 2003, 50(6): 739-745
- [13] Nguyen H, Shindo H. Effects of different levels of compost application on amounts and distribution of organic nitrogen forms in soil particle size fractions subjected mainly to double cropping. Agricultural Sciences, 2011, 2(3): 213-219
- [14] Xin X L, Qin S H, Zhang J B, et al. Yield, phosphorus use efficiency and balance response to substituting long-term chemical fertilizer use with organic manure in a wheat-maize system. Field Crops Research, 2017, 208: 27-33
- [15] 中国科学院南京土壤研究所. 土壤理化分析. 上海: 上海科学技术出版社, 1978: 514-518

- Institute of Soil Science, Chinese Academy of Sciences. Physical and chemical analysis of soil ( In Chinese). Shanghai: Shanghai Science and Technology Press, 1978: 514-518
- [16] Bremner J M. Organic forms of nitrogen // Black C A. ed. Methods of soil analysis. Madison, USA: American Society of Agronomy. 1965:1238-1255
- [17] 鲁如坤. 土壤农业化学分析方法. 北京: 中国农业科学技术出版社,1999: 106-195  
Lu R K. Analytical methods for soil and agricultural chemistry (In Chinese). Beijing: China Agricultural Science and Technology Press, 1999: 106-195
- [18] Chen X, Li Z, Liu M, et al. Microbial community and functional diversity associated with different aggregate fractions of a paddy soil fertilized with organic manure and/or NPK fertilizer for 20 years. *Journal of Soils Sediments*, 2015, 15: 292-301
- [19] Jastrow J D, Amonette J E, Bailey V L. Mechanisms controlling soil carbon turn-over and their potential application for enhancing carbon sequestration. *Climate Change*, 2007, 80: 5-23
- [20] Wang Y D, Wang Z L, Zhang Q Z. Long-term effects of nitrogen fertilization on aggregation and localization of carbon, nitrogen and microbial activities in soil. *Science of the Total Environment*, 2018, 624:1131-1139
- [21] Nieder R, Benbi D K, Scherer H W. Fixation and defixation of ammonium in soils: A review. *Biology and Fertility of Soils*, 2011, 47(1): 1-14
- [22] 丛耀辉. 黑土区水稻土水稳性团聚体有机氮组分及其对矿化氮的贡献. 沈阳: 沈阳农业学, 2016  
Cong Y H. Soil organic nitrogen forms and their contributions to mineralizable nitrogen in water-stable aggregates of paddy soil in black soil region (In Chinese). Shenyang: Shenyang Agricultural University, 2016
- [23] Yang Z P, Zheng S X, Nie J, et al. Effects of long-term winter planted green manure on distribution and storage of organic carbon and nitrogen in water-stable aggregates of reddish paddy soil under a double-rice cropping system. *Journal of Integrative Agriculture*, 2014, 13(8): 1772-1781
- [24] 刘中良, 宇万太, 周桦, 等. 不同有机氮肥输入量对土壤团聚体有机碳组分的影响. *土壤学报*, 2011, 48(6): 1149-1157  
Liu Z L, Yu W T, Zhou H, et al. Effect of application rate of barnyard manure on organic carbon fraction of soil aggregates (In Chinese). *Acta Pedologica Sinica*, 2011, 48(6): 1149-1157
- [25] Schulten H R, Schnitzer, M. The chemistry of soil organic nitrogen: A Review. *Biology and Fertility of Soils*, 1998, 26(1): 1-15
- [26] 张玉树, 丁洪, 王飞, 等. 长期施用不同肥料的土壤有机氮组分变化特征. *农业环境科学学报*, 2014, 33(10): 1981-1986  
Zhang Y S, Ding H, Wang F, et al. Characteristics of organic nitrogen fractions in soils under long-term different fertilization (In Chinese). *Journal of Agro-Environment Science*, 2014, 33(10): 1981-1986
- [27] 丛耀辉, 张玉玲, 张玉龙, 等. 黑土区水稻土有机氮组分及其对可矿化氮的贡献. *土壤学报*, 2016, 53(2): 457-467  
Cong Y H, Zhang Y L, Zhang Y L, et al. Soil organic nitrogen components and their contributions to mineralizable nitrogen in paddy soil of the black soil region (In Chinese). *Acta Pedologica Sinica*, 2016, 53(2): 457-467
- [28] 任金凤, 周桦, 马强, 等. 长期施肥对潮棕壤有机氮组分的影响. *应用生态学报*, 2017, 28(5): 1661-1667  
Ren J F, Zhou H, Ma Q, et al. Effects of long-term fertilization on organic nitrogen fractions in aquatic brown soil (In Chinese). *Chinese Journal of Applied Ecology*, 2017, 28(5): 1661- 1667
- [29] 骆坤, 胡荣桂, 张文菊, 等. 黑土有机碳、氮及其活性对长期施肥的响应. *环境科学*, 2013, 34(2): 676-684  
Luo K, Hu R G, Zhang W J, et al. Response of black soil organic carbon, nitrogen and its availability to



- long-term fertilization (In Chinese). *Environmental Science*, 2013, 34(2): 676-684
- [30] 刘振东, 李贵春, 周颖, 等. 无机肥配施粪肥对华北褐土团聚体分布及有机碳含量的影响. *农业环境科学学报*, 2013, 32(11): 2239-2245
- Liu Z D, Li G C, Zhou Y, et al. The effect of fertilizer management practices on distribution of aggregates and SOC (In Chinese). *Journal of Agro-Environment Science*, 2013, 32(11): 2239-2245
- [31] Ding X L, Liang C, Zhang B, et al. Higher rates of manure application lead to greater accumulation of both fungal and bacterial residues in macroaggregates of a clay soil. *Soil Biology and Biochemistry*, 2015, 84: 137-146
- [32] Andruschkewitsch R, Geisseler D, Dultz S, et al. Rate of soil-aggregate formation under different organic matter amendments—A short-term incubation experiment. *Journal of Plant Nutrition and Soil Science*, 2014, 177: 297-306
- [33] 徐阳春, 沈其荣, 茆泽圣. 长期施用有机肥对土壤及不同粒级中酸解有机氮含量与分配的影响. *中国农业科学*, 2002, 35(4): 403-409
- Xu Y C, Shen Q R, Mao Z S. Contents and distribution of forms of organic N in soil and particle size fractions after long-term fertilization (In Chinese). *Scientia Agricultura Sinica*, 2002, 35(4): 403-409

## Characteristics of the Fraction of Organic Nitrogen in Fluvo-aquic Soil Aggregates under Long-term Application of Chemical Fertilizer and Organic Manure

LI Jiao<sup>1,2</sup> XIN Xiuli<sup>1</sup> ZHU Anning<sup>1†</sup> YANG Wenliang<sup>1</sup> HOU Ying<sup>3</sup> ZOU Wenxuan<sup>3</sup> XU  
Linya<sup>3</sup>

(1 *Institute of Soil Science, Chinese Academy of Sciences/State Key Laboratory of Soil and Sustainable Agriculture/State  
Experimental Station of Agro-Ecosystem in Fengqiu, Nanjing 210008, China*)

(2 *University of Chinese Academy of Sciences, Beijing 100049, China*)

(3 *College of Resources and Environmental Sciences, Nanjing Agricultural University, Nanjing 210095, China*)

**Abstract 【Objective】** The objective of this research was to investigate effects of long-term fertilization on formation of soil aggregates and the fraction of soil organic nitrogen therein in fluvo-aquic soil in the North China Plain, and consequently nitrogen supplying capacity of the soil and its mechanism. **【Method】** Undisturbed soil samples were collected from the four treatments, i.e. CK (no fertilization), NPK (Application of chemical fertilizers only), 1/2OM (Application of half the rate of chemical fertilizer in Treatment NPK and half of the rate of organic manure in Treatment OM) and OM (Application of organic manure only) of the long-term (27 years) fertilization experiment at the State Agro-Ecological Experimental Station in Fengqiu, for analysis of contents of mechanically-stable soil aggregates and organic nitrogen therein, with the dry-sieving method and the Bremner method, separately. **【Result】** Results show that application of organic compost increased the proportion of > 2 mm aggregates in the topsoil, significantly or by 33% and 17% as compared with CK and NPK, whereas it decreased the content of <0.25 mm aggregates. Long-term fertilization significantly increased the content of organic nitrogen in aggregates in the fluvo-aquic soil, especially Treatment OM. The content of organic nitrogen in

aggregates >2 mm, 0.25~2 mm and <0.25 mm was 776.4 mg kg<sup>-1</sup>, 837.7 mg kg<sup>-1</sup> and 625.3 mg kg<sup>-1</sup>, respectively. The organic nitrogen in soil aggregates was dominated with acidolizable ammonium N, which was followed by amino acid nitrogen and unknown-acidolizable nitrogen, aminosaccharide nitrogen, the least. Treatment NPK increased the proportion of acidolizable nitrogen in > 2mm aggregates, while Treatment OM did the content and proportion of amino acid nitrogen and unknown acidolizable nitrogen. 【Conclusion】 long-term application of organic manure can improve the soil structure of fluvo-aquic soil significantly, and stimulate accumulation of total nitrogen and all fractions of organic nitrogen, with amino acid nitrogen, amino sugar nitrogen and non-acidolizable organic nitrogen accumulated mainly in 0.25~2 mm, and acidolizable and unknown acidolizable nitrogen in >2 mm aggregates, thus significantly increasing nitrogen supplying capacity of the soil.

**Key words** Long-term fertilization; Fluvo-aquic soil; Soil aggregates; Acidolizable organic nitrogen; Total nitrogen

(责任编辑: 陈荣府)